

Literatur

- Aldinger, E., Schreiner, M., Bantle, P. (1996):** Standort und Sturmwurf 1990 im östlichen Odenwald. *Agrarforschung in Baden-Württemberg* Bd. 26, 120-129
- Becker, T., Schröter, H., Weigerstorfer, D., Veit, H., editors. (2004):** Waldschutzprobleme nach „Lothar“. Sturm „Lothar“ - Bewältigung der Sturmschäden in den Wäldern Ba.-Wü. LFV BW, Stuttgart.
- Blennow, K., Sallnäs, O. (2004):** WINDA A system of models for assessing the probability of wind damage to forest stands within a landscape. *Ecological Modelling*, 175(1): 87-99.
- FVA (2003):** Dokumentation des Orkanereignisses „Lothar“ sowie Auswertung und Evaluierung der Sturmschadensbewältigung. FVA, Freiburg. 369 S.
- Fridman, J., Valinger, E. (1998):** Modelling probability of snow and wind damage using tree, stand, and site characteristics from *Pinus sylvestris* sample plots. *Scand. J. For. Res.*, 348-356.
- Gardiner, B.A., Quine, C.P. (2000):** Management of forests to reduce the risk of abiotic damage – a review with particular reference to the effects of strong winds. *For. Ecol. Manage.* 135: 261- 277.
- Gardiner, B.A., Peltola, H., Kellomäki, S. (2000):** Comparison of two models for predicting the critical wind speeds required to damage coniferous trees. *Ecological Modelling* 29, 1-23.
- Hanewinkel, M., Zhou, W., Schill, C. (2004):** A neural network approach to identify forest stands susceptible to wind damage. *Forest Ecology and Management* 196: 227-243.
- Hanewinkel, M., Breidenbach, J., Neeff, T. (2008):** 77 years of natural disturbances in a mountain forest area – the influence of storm, snow and insect damage analysed with a long-term time-series. *Canadian Journal of Forest Research* (in review).
- Hinrichs, A. (1994):** Geograph. Informationssysteme als Hilfsmittel der forstl. Betriebsführung. Dis.. Albert-Ludwigs-Univ., Freiburg.
- Jalkanen, A., Mattila, U. (2000):** Logistic regression models for wind and snow damage in northern Finland based on the National Forest Inventory data. *For. Ecol. Manage.* 135: 315-330.
- König, A. (1995):** Sturmgefährdung von Beständen im Altersklassenwald. J.D. Sauerländer's, Frankfurt.
- Kohnle, U., Schmidt, M., Bayer, J., Kändler, G. (2008):** Ein einzelbaumspezifisches Sturmschadensmodell: Grundlagen, Ergebnisse, Anwendung. *FVA einblick+* . 01/08: 24-27.

- Kurth, H., Gerold, D., Dittrich, K. (1987):** Reale Waldentwicklung und Zielwald - Grundlagen nachhaltiger Systemregelung des Waldes, *Wiss. Z. Tech. Univ. Dresd.*, 36, 121-137.
- Lekes, V., Dandul, I. (2000):** Using airflow modelling and spatial analysis for defining wind damage risk classification (WINDARC). *For. Ecol. Manage.* 135, 331-344.
- Mayer, P., Dobbertin, M., Hallenbarter, D., Renaud, J.-P., Walthert, L., Zimmermann, S. (2005):** Forest storm damage is more frequent on acidic soils. *Annals of Forest Science* 62:303-311.
- Miller, D.R., Dunham, R., Broadgate, M.L., Aspinall, R.J., Law, A.N.R. (2000):** A demonstrator of models for assessing wind, snow and fire damage to forests using the WWW. *For. Ecol. Manage.* 135, 355-363.
- Mitchell, S.J., Hailemariam, T., Kulis, Y. (2001):** Empirical modelling of cutblock edge windthrow risk on Vancouver Island, Canada, using stand level information. *For. Ecol. Manage.* 154, 117-130.
- Peltola, H., Kellomäki, S., Väisänen, H., Ikonen, V.-P. (1999):** A mechanistic model for assessing the risk of wind and snow damage to single trees and stands of Scots pine, Norway spruce, and birch. *Can. J. For. Res.* 29, 647-661.
- Rottmann, M. (1985):** Schneebruchschäden in Nadelholzbeständen. J.D. Sauerländer's, Frankfurt a.M.
- Rottmann, M. (1986):** Wind- und Sturmschäden im Wald. J.D. Sauerländer's, Frankfurt a.M.
- Schelhaas, M.-J., Nabuurs, G.-J., Schuck, A. (2003):** Natural disturbances in the European forests in the 19th and 20th centuries. *Glob. Chang. Biol.* 9: 1620-1633.
- Schütz, J. P., Götz, M., Schmid, W., Mandallaz, D. (2006):** Vulnerability of spruce (*Picea abies*) and beech (*Fagus sylvatica*) forest stands to storms and consequences for silviculture. *Eur. J. For. Res.* 125: 291-302.
- Suzuki, T. (1971):** Forest transition as a stochastic process. *Mitt. Forstl. Bundes-Vers.anst. Wien* 91, 137-150.
- Talkkari, A., Peltola, H., Kellomäki, S., Strandmann, H. (2000):** Integration of component models from the tree, stand and regional levels to assess the risk of wind damage at forest margins. *For. Ecol. Manage.* 135, 303-313.
- Valinger, E., Fridman, J. (1997):** Modelling probability of snow and wind damage in Scots pine stands using tree characteristics. *For. Ecol. Manage.* 97, 215-222.
- Valinger, E., Fridman, J. (1999):** Models to assess the risk of snow and wind damage in pine, spruce, and birch forests in Sweden. *Environ. Manage.* 24, 209-217.